

Itinerant ferromagnetism in an atom trap

Ilya Berdnikov,^{1,2} P. Coleman,¹ and Steven H. Simon²

¹Center for Materials Theory, Rutgers University, Piscataway, New Jersey 08854, USA

²Alcatel-Lucent, Bell Labs, Murray Hill, New Jersey 07974, USA

(Received 12 April 2009; published 2 June 2009)

We propose an experiment to explore the magnetic phase transitions in interacting fermionic Hubbard systems and describe how to obtain the ferromagnetic phase diagram of itinerant-electron systems from these observations. In addition, signatures of ferromagnetic correlations in the observed ground states are found: for large trap radii (trap radius $R_T > 4$, in units of coherence length ξ), ground states are topological in nature—a “skyrmion” in two dimension, and a “hedgehog” in three dimension.

DOI: [10.1103/PhysRevB.79.224403](https://doi.org/10.1103/PhysRevB.79.224403)

PACS number(s): 67.85.-d, 71.10.Fd

I. INTRODUCTION

The simplest and best studied model of itinerant ferromagnetism (FM) is the Hubbard model.¹ Shortly after its introduction, Nagaoka and Thouless proved^{2,3} that for an interaction U of infinite strength, doping one hole into a background of spins leads to FM. Yet for more than forty years, the fate of the Thouless-Nagaoka phase in the Hubbard model at finite doping and finite interaction has not been fully resolved. Existing studies include various perturbative,^{4–6} variational,^{7–10} slave boson,^{11,12} Quantum Monte Carlo,^{13,14} and DMFT (Ref. 15) calculations. For example, there is no consensus as to the values of the critical doping δ_{cr} ($\approx 0.19–0.49$) below which FM occurs and critical interaction U_{cr} ($\approx 63–77.7$ in units of the hopping amplitude t). Moreover, the various approaches do not agree on the nature of the transition, first order claimed by some^{4,5,11} second order by others.^{13–15}

Experiments on optical lattices, which allow a tunable control of model parameters, offer an interesting opportunity to address these long-standing open questions. This setting was used to study correlations in cold bosonic systems experimentally^{16,17} and could also be applied to superconductivity of fermions. The superconducting transition, however, requires a low entropy state cooled to temperatures far below degeneracy, which still poses a significant challenge. Ferromagnetism offers a particular advantage in this respect, since, at least for the Nagaoka phase at $U=\infty$, the absence of any other scale sets the transition temperature to be a finite fraction of the bandwidth, and entropy at the transition is not a small parameter (of order unity per carrier). A recent experiment on two-component fermionic ⁴⁰K in optical lattices has reported a Mott insulator with a maximum achieved $U/t \sim 40$, $T/T_F = 0.28$, and initial densities of more than one electron per site.¹⁸ A FM phase presumably resides near the observed Mott phase and conservative estimates place it at slightly less than one particle per site, $U/t \sim 100$ and $T/T_F \sim 0.1$.^{9,10,15} The experimental parameters required in a setup such as¹⁸ are estimated: lattice depth $V_0/E_r \sim 13–19$, depending on the scattering length a_s , the coherence length $\xi = \frac{\hbar}{\sqrt{2m\mu}} \sim 0.2 \mu\text{m}$, and the trap radius $R_T \sim 1.6 \mu\text{m}$, obtained from the laser waist radius of $160 \mu\text{m}$.

In this paper we explore the possibility of using cold fermion gases to study FM in both two-dimensional (2D) and

three-dimensional (3D) systems. For large trap sizes (radius $R_T > 4$ and coherence lengths ξ) with the constraint of zero total magnetization and filling factor less than unity everywhere in the trap, we find that the “skyrmion” configuration [Fig. 1(a)] has the lowest energy in 2D while in 3D the lowest energy configuration is the “hedgehog” [Fig. 1(b)]. Observing these ground states provides a simple way to map out the FM phase diagram which would be the first result of its kind.

II. APPROACH

Our approach to the problem is in marked contrast with previous studies,^{19–22} some of which address only the 2D case: rather than using cold atoms as a means to verify microscopic calculations, we propose using them as a direct probe of itinerant FM in both 2D and 3D traps. Moreover, in the analysis in Ref. 20 total magnetization is not conserved; whereas, in most of the recent experiments,^{23–25} total magnetization is, in fact, conserved, as a consequence of isolation from the environment and an absence of coupling between the effective spin degree of freedom and the rest of the system.

Ferromagnetism has been studied extensively in the context of multicomponent Bose-Einstein condensates (spinor BECs), primarily in two dimensions, starting with Refs. 26–31. However, magnetic order in these systems is a secondary consequence of condensation and develops wherever the condensate forms. Fermions near the point of degeneracy, on the other hand, become FM in the absence of any other broken symmetry. Trapping potentials give rise to spatial density variations, so different regions of the fermionic fluid at or near degeneracy may not reach the Stoner instability³²




FIG. 1. (Color online) (a) Skyrmion and (b) hedgehog configuration of spins.

simultaneously, resulting in separation between the FM and paramagnetic (PM) phases,¹⁹ something intrinsically absent in spinor BECs and bulk materials. This phase separation ensures that the resulting ferromagnetic droplets have definite size, and this, in turn, allows one to measure the phase diagram, as discussed below.

We consider an optical lattice containing fermionic atoms cooled to temperatures close to degeneracy. The role of spin is played by two states in the atoms' hyperfine multiplet. Hopping between lattice sites is determined by the overlap of the atomic wave functions on those sites, while the on-site interaction term is given by the s -wave scattering of "electrons" with opposite "spin."³³ In a trapped gas of equal spin populations, ground states are constrained by vanishing magnetization $\vec{M} = \int d\mathbf{r} \vec{M}(\mathbf{r}) = 0$.

What kind of ferromagnetic quantum state satisfies this condition? One possibility is the state $|\psi\rangle = |S, 0\rangle$ with maximal total $\vec{M}^2 = S(S+1)$ but $M_z = 0$. The expectation values of M_x and M_y in this state are zero, ostensibly satisfying the requirement that the total magnetization $\langle \psi | \vec{M} | \psi \rangle = 0$. However, we can rule out such a state by observing that each component of the total magnetization operator is conserved, $[H, M_\alpha] = 0$, so that the quadrupole operator $Q_{\alpha\beta} = M_\alpha M_\beta - \frac{1}{3} M^2 \delta_{\alpha\beta}$ is also conserved: $[H, Q_{\alpha\beta}] = 0$. The state $|\psi\rangle$ has a nonvanishing quadrupole moment, where $\langle Q_{xx} \rangle = \langle Q_{yy} \rangle = -2 \langle Q_{zz} \rangle = S(S+1)/6$, and therefore cannot develop as a spin-isotropic paramagnetic state is cooled through the ferromagnetic transition.

III. LANDAU-GINZBURG TREATMENT

To go further, we need to consider states of nonuniform magnetization in which $\vec{M} = 0$. To this end, we make a long-wavelength expansion of the total energy as a functional of the local magnetization. Such a long-wavelength treatment of the problem does not imply that the underlying nature of the system is classical. Indeed, long-wavelength actions of this sort have been used to great success in the analysis of one-dimensional quantum antiferromagnets, where long-range order is completely absent.³⁴

The Landau-Ginzburg free-energy functional which describes long-wavelength configurations of the magnetization order parameter takes the form

$$F_{\text{LG}} = \int d\mathbf{r} \left[\frac{\rho}{2} |\nabla \vec{M}|^2 + \frac{\beta}{4} \left(|\vec{M}|^2 + \frac{\alpha(\mathbf{r})}{\beta} \right)^2 \right]. \quad (1)$$

Odd-power terms are ruled out by the time-reversal symmetry of the free energy. Coefficients ρ and β are assumed to be positive and constant for simplicity and the entire effect of the trap potentials is in the position dependence of $\alpha(\mathbf{r})$. We define R_c such that $\alpha(\mathbf{r}) < 0$ for $r < R_c$ and $\alpha(\mathbf{r}) > 0$ for $r > R_c$. We assume that the density everywhere in the system is less than one electron per site and conclude with a discussion of the remaining cases.

Clearly, whatever magnetic moment is accumulated by any one region in the trap must be completely canceled by the rest of the trap. The configuration which connects any two regions could either be a domain wall or some form of a




FIG. 2. (Color online) Showing (a) domain wall and (b) single-axis twist configuration of spins 3.

twist (either with or without vorticity). It is the competition between these three scenarios that determines the ground state. Due to global rotation invariance, it suffices to consider a single representative state from each class.

In 3D the first candidate is a hedgehog [Fig. 1(b)]. Qualitatively, it consists of a core of radius r_0 , where magnetization is suppressed, a region of thickness L in which it continuously increases from 0 to some fraction $|a| \leq 1$ of the uniform value M_0 , and the outer region, extending to the edge of the trap at radius $R_T = R_c$. In a particular realization, the magnetization vector at any point is in the radial direction. A competing configuration has a core of radius r_0 maximally polarized in one direction. A domain wall of thickness L connects the core to the outer region polarized at a fraction $0 \leq a \leq 1$ of the maximum M_0 in the opposite direction [Fig. 2(a)]. The remaining possibility is a pure twist. The twist plane is globally fixed (a rotation of θ_0 about the z axis) and the magnetization turns about the x axis through an angle $a\pi$, $1 \leq a \leq 2$. The twist occupies a shell of thickness L outside of a maximally polarized core of radius r_0 and the remaining outer region is polarized along the final direction [Fig. 2(b)]. Parameters a , r_0 , and L for each configuration are determined by numerically minimizing F_{LG} for a fixed value of R_c subject to the constraint of zero net magnetization.

In 2D in addition to the domain wall, the single-axis twist and the hedgehog, which are just planar slices of their 3D cousins, there is another possibility: the skyrmion. This configuration has a maximally polarized core of radius r_0 at the center, a twist through the angle $a\pi$, $1/2 \leq a \leq 1$ in a ring of thickness L , and an outer region, polarized along the final direction [Fig. 1(a)]. Unlike the single-axis twist, the twist axis of the skyrmion is a function of position. The constraint for the 2D candidates must be implemented explicitly and the ground state is obtained in the same way as in 3D.

Before solving the full problem, some estimates are in order. In 2D a domain wall is approximately the suppression of the order parameter in an annulus of thickness L and inner radius r_0 . Omitting a dimensionful prefactor common to all configurations, and in the case of the domain wall, ignoring the stiffness contribution, we find the free energy ($\xi = \sqrt{\rho/\alpha}$ is the coherence length)

$$F_{\text{DW}} \sim \frac{2r_0L + L^2}{2\xi^2}. \quad (2)$$

The energy is minimized by shrinking L to $L \propto \xi$, however, the magnetization constraint forces $r_0 \propto R_c$, and so $F_{\text{DW}} \sim R_c$.




FIG. 3. (Color online) (a) Free energy vs trap size in two dimensions, ● hedgehog, ▲ skyrmion, ◆ domain wall, and ■ single-axis twist. Insert: $\text{Log}_e(F)$ vs R_c/ξ . (b) Free energy vs size of the correlated region in three dimensions for a fixed trap size, $R_T=75\xi$, ● hedgehog, ◆ domain wall, and ■ single-axis twist. Insert: $\text{Log}_e(F)$ vs R_c/ξ .

In the same units the skyrmion free energy has contributions from vorticity and from twisting, i.e.,

$$F_{\text{SK}} \sim 2 \ln \frac{R_c}{r_0} + (a\pi)^2 \left(\frac{r_0}{L} + \frac{1}{2} \right). \quad (3)$$

The total magnetic moment of the skyrmion is zero for some $a \sim O(1)$. F_{SK} is minimized when both $L, r_0 \propto R_c$. As a result, the skyrmion free energy is roughly independent of the trap size. Analysis of the single-axis twist is similar, however, implementing the constraint requires more twisting in the same volume than in the skyrmion case, and therefore this configuration should always be higher in energy. The 2D hedgehog free energy is estimated to be $F_{\text{Hg}} \sim 2 \ln \frac{R_c}{r_0} + \frac{1}{2} \left(\frac{r_0}{\xi} \right)^2$, sum of vorticity and core energy, and one can show that $F_{\text{Hg}} \sim \ln R_c/\xi$. Thus, for large trap sizes we expect a skyrmion to form in the trap. Similar treatment of the 3D configurations shows that the hedgehog should be the ground state. Numerical computations confirm these estimates, as shown in Fig. 3. We find that in small traps in both 2D and 3D domain walls are preferred, while for large traps in 3D ($R_c/\xi > 7$) the hedgehog has the lowest energy

and in 2D ($R_c/\xi > 4$) the skyrmion is the ground state.

In the above derivation we tacitly assumed that the whole system becomes FM, i.e., $R_c=R_T$. This is close to the truth in 2D since the density of states is practically flat and the Stoner criterion for magnetic instability dictates that the entire system becomes FM at once independent of the spatial density variations. In 3D, however, the density of states is more complicated, thus certain parts of the trap will cross into the broken-symmetry regime earlier than others. In the language of Landau-Ginzburg, it means that α changes sign from positive outside of some radius R_c , to negative inside. Incorporating this into our analysis requires somehow suppressing the magnetization outside R_c in the candidate states. This can be accomplished by half of a domain wall, e.g., $\vec{M}(r)$ diminishing to 0 within some shell of thickness L_1 of the critical radius R_c and remaining 0 to the edge of the trap R_T . Although the qualitative results of the calculations remain the same, the numerical problem itself is modified: we minimize F_{LG} with respect to parameters a, r_0, L , and L_1 for a given radius of the FM region (R_c), and a given radius of the trap (R_T). Results for 3D in Fig. 3 properly reflect these considerations.

In the case when the trap has regions with density higher than one electron per site, we expect further phase separation. In these regions the chemical potential may enter the Mott gap and we would expect an antiferromagnetic Mott phase to set in with filling locked to one electron per site. In regions of even higher density, the chemical potential might cross the gap and emerge, once again, in an itinerant band. Thus, we expect the phase profile to have wedding-cake structure.¹⁷ For traps with very high density in the middle, we expect a hole PM at the core, followed in turn by shells of hole FM, an antiferromagnetic Mott insulator, and an electron FM, finally ending with the electron PM exterior. For intermediate densities, the Mott insulator could form a natural core for the 3D hedgehog configurations.

IV. DISCUSSION

Experiments with ultracold atomic gases generally exploit absorption imaging to detect the state of matter in the trap. For multicomponent gases *in situ* imaging of each individual component is possible.^{22,24,25,35} In the case of FM systems this technique allows resolving the full magnetization integrated along the camera axis. Figure 4 shows *in situ* images of integrated model magnetization for the hedgehog and the skyrmion. The critical radius can be clearly seen (and measured) in the 3D configuration and it designates the phase boundary between the PM outside and the FM inside. In addition to the phase boundary and the profile of the magnetization, one could look for signatures of correlation in the shot noise.^{36,37} The autocorrelator $\langle M_z(\mathbf{r})M_z(\mathbf{r}') \rangle$ can be calculated using the time-of-flight images since it is related to an integral of the frequency-space-magnetization correlator. Tracking the associated correlation length, which will diverge as the system approaches the transition, is, potentially, a more sensitive tool and should indicate the proximity of the FM phase before the image itself shows the phase separation.³¹




FIG. 4. (Color online) False color plot of magnetization integrated *in situ*, symmetry broken along the z axis. Hedgehog imaged along the x axis (left) and skyrmion imaged along the z axis (right).

With the newly found signatures of correlation we can map out the phase diagram of an interacting fermion gas. The density of particles at the critical radius R_c can be obtained from the absorption images. Properly normalized, e.g., the Mott insulator is exactly at half-filling, this density defines the critical doping of the system. Interaction strength $U/t \sim a_s \exp(2\sqrt{V_0/E_f})$ is given by the s -wave scattering length, a_s , and the lattice depth V_0 with a_s and V_0 (i.e., U and t) controlled independently.³³ Thus scanning a_s and V_0 and measuring the critical doping for each, we determine the phase diagram. As discussed above, at higher densities the magnetization images should exhibit an even richer shell structure. However, by particle-hole symmetry all of the FM regions should have the same basic features and each phase boundary present in the image will provide a point on the full phase diagram. The finer details of the observed configuration can give us even more details about the parameters of the effective Landau-Ginzburg description; and in turn our observation of the Landau-Ginzburg parameters can be used to evaluate the efficacy of various microscopic calculational schemes. Similarly, we could explore the vicinity of the FM quantum critical point and compare the observations to the Hertz-Millis theory. In any real experiment a number of practical issues will play a role. In particular, the details of symmetry breaking must be the same in successive experimental

runs and in the different planes of a quasi-2D-trap configuration. One way to ensure this is to impose a small position-dependent external field. A weak interplanar coupling in quasi-2D lattice might also stabilize the long-range order in the trap. Our model neglects the fact that in experiments to date, strong interactions mix the various bands of the optical lattice.^{38,39} Elimination of this effect may require a lower interaction strength, which in turn will require a still lower temperature.³³

In this paper we demonstrated that ground states of interacting fermions in both 2D and 3D traps are configurations with nontrivial topologies. We have determined detectable signatures of these states and proposed an experiment to map the phase diagram using these signatures. Such experiments offer the prospect of directly probing itinerant FM, studying the Stoner criterion in strongly correlated magnets, and providing much needed experimental input in the debate on the phase diagram of the Hubbard model.

ACKNOWLEDGMENTS

The authors would like to thank T.-L. Ho and T. Esslinger for fruitful discussions. This work was supported by NSF under Grant No. DMR-0605935, DOE under Grant No. DE-FE02-00ER45790, and the Rutgers-Lucent Foundation.

¹J. Hubbard, Proc. R. Soc. London, Ser. A **276**, 238 (1963).

²Y. Nagaoka, Phys. Rev. **147**, 392 (1966).

³D. J. Thouless, Proc. Phys. Soc. London **86**, 893 (1965).

⁴A. Oles, J. Phys. C **15**, L1065 (1982).

⁵L. B. Ioffe and A. I. Larkin, Phys. Rev. B **37**, 5730 (1988).

⁶W. O. Putikka, M. U. Luchini, and M. Ogata, Phys. Rev. Lett. **69**, 2288 (1992).

⁷B. S. Shastry, H. R. Krishnamurthy, and P. W. Anderson, Phys.

Rev. B **41**, 2375 (1990).

⁸W. von der Linden and D. M. Edwards, J. Phys.: Condens. Matter **3**, 4917 (1991).

⁹T. Hanisch and E. Müller-Hartmann, Ann. Phys. **505**, 381 (1993).

¹⁰P. Wurth, G. Uhrig, and E. Müller-Hartmann, Ann. Phys. **5**, 148 (1996).

¹¹B. Möller, K. Doll, and R. Frésard, J. Phys.: Condens. Matter **5**,

- 4847 (1993).
- ¹²D. Boies, F. A. Jackson, and A.-M. S. Tremblay, *Int. J. Mod. Phys. B* **9**, 1001 (1995).
- ¹³X. Y. Zhang, E. Abrahams, and G. Kotliar, *Phys. Rev. Lett.* **66**, 1236 (1991).
- ¹⁴F. Becca and S. Sorella, *Phys. Rev. Lett.* **86**, 3396 (2001).
- ¹⁵H. Park, K. Haule, C. A. Marianetti, and G. Kotliar, *Phys. Rev. B* **77**, 035107 (2008).
- ¹⁶M. Greiner, O. Mandel, T. Esslinger, T. W. Hänsch, and I. Bloch, *Nature (London)* **415**, 39 (2002).
- ¹⁷R. B. Diener, Q. Zhou, H. Zhai, and T.-L. Ho, *Phys. Rev. Lett.* **98**, 180404 (2007).
- ¹⁸R. Jördens, N. Strohmaier, K. Gnter, H. Moritz, and T. Esslinger, *Nature (London)* **455**, 204 (2008).
- ¹⁹L. Salasnich, B. Pozzi, A. Parola, and L. Reatto, *J. Phys. B* **33**, 3943 (2000).
- ²⁰T. Sogo and H. Yabu, *Phys. Rev. A* **66**, 043611 (2002).
- ²¹R. A. Duine and A. H. MacDonald, *Phys. Rev. Lett.* **95**, 230403 (2005).
- ²²Shizong Zhang, Hsiang-hsuan Hung, and Congjun Wu, arXiv:0805.3031 (unpublished).
- ²³M. Zwierlein, A. Schirotzek, C. H. Schunck, and W. Ketterle, *Science* **311**, 492 (2006).
- ²⁴G. B. Partridge, W. Li, R. I. Kamar, Yean-an Liao, and Randall G. Hulet, *Science* **311**, 503 (2006).
- ²⁵Y. Shin, M. W. Zwierlein, C. H. Schunck, A. Schirotzek, and W. Ketterle, *Phys. Rev. Lett.* **97**, 030401 (2006).
- ²⁶T.-L. Ho, *Phys. Rev. Lett.* **81**, 742 (1998).
- ²⁷T. Ohmi and K. Machida, *J. Phys. Soc. Jpn.* **67**, 1822 (1998).
- ²⁸R. A. Battye, N. R. Cooper, and P. M. Sutcliffe, *Phys. Rev. Lett.* **88**, 080401 (2002).
- ²⁹H. Saito and M. Ueda, *Phys. Rev. A* **72**, 023610 (2005).
- ³⁰A. E. Leanhardt, Y. Shin, D. Kielpinski, D. E. Pritchard, and W. Ketterle, *Phys. Rev. Lett.* **90**, 140403 (2003).
- ³¹L. E. Sadler, J. M. Higbie, S. R. Leslie, M. Vengalattore, and D. M. Stampfer-Kurn, *Nature (London)* **443**, 312 (2006).
- ³²E. C. Stoner, *Proc. R. Soc. London, Ser. A* **165**, 372 (1938).
- ³³I. Bloch, J. Dalibard, and W. Zwerger, *Rev. Mod. Phys.* **80**, 885 (2008).
- ³⁴F. D. M. Haldane, *Phys. Rev. Lett.* **50**, 1153 (1983).
- ³⁵Y. Shin and C. H. Schunck, A. Schirotzek, and W. Ketterle, *Nature (London)* **451**, 689 (2008).
- ³⁶E. Altman, E. Demler, and M. D. Lukin, *Phys. Rev. A* **70**, 013603 (2004).
- ³⁷T. Rom, Th. Best, D. Van Oosten, U. Schneider, S. Göllin, B. Paredes, and I. Bloch, *Nature (London)* **444**, 733 (2006).
- ³⁸M. Kohl, H. Moritz, T. Stoferle, K. Gunter, and T. Esslinger, *Phys. Rev. Lett.* **94**, 080403 (2005).
- ³⁹R. B. Diener and T.-L. Ho, *Phys. Rev. Lett.* **96**, 010402 (2006).